

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 208 375
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 86201160.8

(51) Int. Cl.4: A61K 39/116 , A61K 39/385 ,
C07K 15/14

(22) Date of filing: 02.07.86

(30) Priority: 05.07.85 IT 2145185

(43) Date of publication of application:
14.01.87 Bulletin 87/03

(64) Designated Contracting States:
AT BE CH DE FR GB LI LU NL SE

(71) Applicant: SCLAVO S.p.A.
Via Fiorentina 1
I-53100 Siena(IT)


(72) Inventor: Porro, Massimo
Via S. Marco 43
I-56010 Localita' Collanza Asciano Siena(IT)
Inventor: Costantino, Paolo
Via Toscana 11
I-53034 Colle Di Val D'Elsa(Siena(IT)

(74) Representative: Roggero, Sergio et al
Ing. Barzanò & Zanardo Milano S.p.A. Via
Borgonuovo 10
I-20121 Milano(IT)

(54) Glycoprotein conjugates having trivalent immunogenic activity.

(57) Glycoprotein conjugates having trivalent immunogenic activity obtained by binding, by a covalent bond, to a protein selected among CRM 197, tetanus toxoid, and pertussis toxin, at least an oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium and at least an oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, and wherein said oligosaccharidic haptens are previously activated by introducing terminal esters.

Fig.1



EP 0 208 375 A2

The present invention relates to glycoprotein conjugates having trivalent immunogenic activity, to their preparation, and to their use as vaccines.

In particular, the present invention relates to glycoprotein conjugates having trivalent immunogenic activity obtained by binding, by a covalent bond, to a protein selected among CRM 197, tetanus toxoid and pertussis toxin, at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, after the preliminary activation of said oligosaccharidic haptens by the introduction of terminal ester groups.

The development of vaccines against such capsulate bacteria as meningococcus and pneumococcus by the use of capsular polysaccharidic antigens derived from said bacteria, has shown the poor immunogenicity of said polysaccharidic antigens, above all in infantile population.

This represents a problem of particular gravity and interest because the greatest level of incidence of infections caused by said capsulate bacteria occurs in children. From this the necessity derives of improving the immunogenic properties of capsular polysaccharides for the preparation of more effective vaccines.

The synthesis and the use of glycoprotein conjugates represent a particularly interesting solution in improving the immune response to the polysaccharidic antigens.

Avery (J. Exp. Med. (1929) 50, 533) and Gcbel (J. Exp. Med. (1939) 69, 53) described, about 50 years ago, the preparation and the use of glycoprotein conjugates, demonstrating that the immunization of animals by the said compounds, wherein the saccharidic moiety was polysaccharide of *S. Pneumoniae* Type 3, determined an immunization against the bacterium, which was more efficacious than that of polysaccharide as such.

Many studies have been recently published in the field of the synthesis of the glycoprotein conjugates, and promising results have been obtained with *Haemophilus influenzae* Type b, and *Meningococcus* Group A and C.

U.S. Pat. 4,356,170 discloses and claims a glycoprotein conjugate constituted by the capsular polysaccharide of an either gram-positive or gram-negative bacterium and by a protein, and a process for the preparation thereof, characterized in that the capsular polysaccharide is modified, before being bound to the protein, so to generate terminal aldehydic groups by oxydation of vicinal hydroxy groups.

According to said technique of the prior art, the glycoprotein conjugates so obtained can be used as vaccines against gram-positive or gram-negative bacteria, or for the preparation of multiple vaccines, i.e., vaccines with immune responses to different bacteria, by mixing two or more glycoconjugates in a pharmaceutically acceptable liquid carrier.

It has been found now that, under particular conditions, binding is possible, to a same protein, oligosaccharidic haptens derived from capsular polysaccharides of gram-negative and gram-positive bacteria, a glycoprotein conjugate having trivalent immunogenic activity being obtained.

Accordingly, the present invention relates to glycoprotein conjugates having trivalent immunogenic activity, obtained by binding, by a covalent bond, to a protein selected among CRM 197, tetanus toxoid, pertussis toxin, at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, after the preliminary activation of the oligosaccharidic haptens by the introduction into them of terminal ester groups.

The glycoprotein conjugates so obtained are capable of inducing on an animal model, contemporaneously, anamnestic immune responses to the capsular polysaccharidic antigen of gram-negative bacteria, to the capsular polysaccharidic antigen of gram-positive bacteria, as well as to the protein used. The conjugates of the present invention are useful to prepare novel vaccines against epidemiologically meaningful capsulate gram-negative and gram-positive bacteria, by using one single glycoprotein conjugate having trivalent immunogenic activity.

According to the present invention, the proteins used for the preparation of the glycoprotein conjugates having trivalent immunogenic activity, are selected among the proteins physiologically tolerated and with free amino groups.

Preferably, proteins having thymus-dependant immunogenic activity are used.

Among these, particularly preferred are CRM 197, tetanus toxoid and pertussis toxin.

Said proteins are prepared by culturing the respective producer microorganisms, separating from the culture medium by filtration, precipitating and finally purifying them by chromatographic techniques.

According to the present invention, the oligosaccharidic haptens are selected among those derived from capsular polysaccharides of epidemiologically meaningful gram-positive and gram-negative bacteria.

Particularly suitable to the purpose are those derived from gram-positive bacteria belonging to the group of *Streptococcus pneumoniae*, *Streptococcus β-emoliticus*, and those derived from capsular polysaccharides of gram-negative bacteria belonging to the group of *Neisseria meningitidis*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Haemophilus influenzae*.

The oligosaccharidic haptens are prepared by acidic hydrolysis of the corresponding capsular polysaccharidic antigens. Typically, the hydrolysis reaction is carried out by suspending the capsular polysaccharide in an aqueous solution of acetic acid at a pH value comprised within the range of from 3 to 5.5.

The hydrolysis is carried out by maintaining the solution inside sealed ampuls, at a temperature of 100°C, for a time of from 6 to 40 hours.

At the end of the hydrolysis reaction, the oligosaccharidic haptens are separated from the reaction mixture by gel-chromatography, and the effluents are analysed to determine the presence of methyl-pentoses, phosphorus and reducing groups.

The molecular weight of said oligosaccharidic haptens is determined by gel-chromatographic analysis, by measuring the distribution constant - (Kd), whilst the characterization of the physical dimensions is carried out by N.M.R. analysis.

The immunochemical properties of said oligosaccharidic haptens are determined according to the method as disclosed by Porro et al. (*Molec. Immunol.*, in press, 1985), which combines the technique of electrophoretic affinity with the "rocket" immunoelectrophoresis technique, and is based on the inhibiting capacity of the haptens in the immunoprecipitation reactions between the antigen and homologous antibodies.

Oligosaccharidic haptens suitable to the purpose are selected among those having a molecular weight of from 10^3 to 2.10^5 , preferably having a molecular weight of 1500, and characterized by the presence of terminal reducing groups.

According to the present invention, said oligosaccharidic haptens are activated by the introduction of terminal ester groups before being bound by covalent bonds to the protein.

The activation comprises the introduction of primary amino groups into the terminal reducing groups of the oligosaccharidic haptens, and the subsequent conversion of said amino groups into the corresponding esters.

The process is generally carried out by suspending the oligosaccharidic haptens in water in the presence of ammonium chloride and NaBH_4CN .

The solution so obtained is adjusted to a pH value of from 7.5 to 9.5 and is kept, within sealed ampuls, at a temperature of from 30°C to 55°C for a corresponding time of from 1 to 2 weeks.

At the end of the reaction, the oligosaccharidic haptens are separated by gel-chromatography and the effluents are analyzed to determine the presence of methylpentoses, phosphorus and amino groups.

The oligosaccharidic haptens so obtained are subsequently transferred into dimethylsulphoxide and esterified in the presence of esters of organic acids, such as derivatives of adipic acid.

Preferably, the disuccinimidyl ester of the hydrazide of adipic acid is used, at a molar concentration in excess relatively to the amino groups introduced into the oligosaccharidic haptens, generally ranging from 2:1 to 10:1.

The oligosaccharidic haptens so activated are precipitated from the reaction mixture by the addition of 1,4-dioxane, using a 1,4-dioxane/dimethylsulphoxide ratio of 10:1 by volume.

The activated oligosaccharidic haptens are then centrifuged off from the reaction mixture, and analyzed to determine the disappearing of the amino groups and the conversion thereof into active esters.

The present invention relates moreover to a process for the preparation of glycoproteic conjugates having trivalent immunogenic activity comprising the binding, contemporaneously or sequentially, by a covalent bond, to a protein selected among CRM 197, tetanus toxoid or pertussis toxin, at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, after the preliminary activation of the oligosaccharidic haptens by the introduction of terminal ester groups.

The reaction of conjugation between the protein and the oligosaccharidic haptens is carried out by adding to a protein, either contemporaneously or sequentially, the oligosaccharidic haptens, previously activated, operating with a protein/hapten molecular ratio of 1:10.

Typically, the conjugation reaction is carried out in the liquid phase, in the presence of an organic solvent, at a temperature comprised within the range of from 10°C to 40°C, for a corresponding time period of from 2 to 24 hours. Preferably, the process is carried out at a temperature of 15°C for a time of 15 hours, in the presence of dimethylsulphoxide.

At the end of the conjugation reaction, from the reaction mixture, the glycoproteic conjugates are separated by gel-chromatography, the chromatographic effluents showing a chemical activity to-

wards the protein and towards the bound oligosaccharidic haptens are combined. The effluents are sterilized by filtration, by using filter membranes of 0.22 μ m.

The substitution degree in the proteinic molecule is verified for the polysaccharidic haptens by the chemical analysis of the glycoproteinic conjugate and subsequent conversion into mol of the percentages of the three components.

According to the present invention, the substitution degree of the glycoproteinic conjugate having trivalent immunogenic activity is of at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium and of at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium per each mol of protein. Generally, the number of oligosaccharidic haptens derived from the capsular polysaccharide of a gram-positive bacterium is comprised within the range of from 1 to 6, and that of the oligosaccharidic haptens derived from the capsular polysaccharide of a gram-negative bacterium of from 1 to 2 per each mol of protein.

The glycoproteinic conjugates of the present invention are useful as vaccines against epidemiologically meaningful capsulate gram-positive bacteria and capsulate gram-negative bacteria.

In particular, they can be used as vaccines to immunize the infantile population. Such molecules contain the oligosaccharidic haptens derived from capsular polysaccharides, the immune response of which is thymus-independent.

Said polysaccharides, by themselves poorly immunogenic in children, when are used for the preparation of the glycoproteinic conjugates of the present invention, yield on animal models a thymus-dependent immune response.

In particular, a test carried out by the ELISA method has demonstrated the production of antisera of antibodies specific to capsular polysaccharides, when the glycoproteinic conjugates of the present invention are used.

According to the present invention, vaccines are prepared by mixing a therapeutically effective amount of the glycoproteinic conjugate having trivalent immunogenic activity, with a liquid carrier selected among a physiologic solution, or other injectable liquid carriers.

To the said mixture, additives selected among those known from the present art for the preparation of vaccines can be added. Additives suitable to that purpose are such stabilizer compounds as lactose or sorbitol, and such adjuvants as aluminum hydroxide, phosphate, or an alginate.

Preferably, AlPO₄ is used.

The vaccines so obtained are administered by intramuscular or subcutaneous way.

The concentration of the glycoproteinic conjugates for the preparation of vaccines for children is comprised within the range of from 25 to 200 μ g. Greater doses can be administered on the basis of the body weight.

The vaccine can be administered as one single dosis, or it can be divided into a plurality of doses.

To the purpose of illustrating the present invention, without however desiring limiting it, a glycoproteinic compound has been synthesized, wherein protein CRM 197 is bound by a covalent bond to at least one oligosaccharidic hapten derived from the capsular polysaccharide of the gram-positive bacterium *Streptococcus pneumoniae* Type 6A, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of the gram-negative bacterium *Neisseria meningitidis* group C.

The immunogenic properties of said glycoproteinic conjugate have been tested on animals, evaluating the immune responses to protein CRM 197, as well as to the capsular polysaccharides of *S. pneumoniae* Type 6A, and of *N. meningitidis* group C.

Description of the Figures

Figure 1: the electrophoretic behaviour is shown of the glycoproteinic conjugate (5 μ g, line 2) and of the native protein CRM 197 (5 μ g, line 3).

The chemical glycosilizing appears distributed on both the A and B subunits forming the protein CRM 197.

The molecular weight of the glycoproteinic conjugate, computed on the basis of the molecular weights shown at line 1, is of about 78,000.

Figure 2: The chart is shown which is obtained by reporting on the abscissae the time (days) and on the ordinates the concentrations of IgG immunoglobulins, expressed as ELISA units.

By —●— the values are shown of anti-*S. pneumoniae* Type 6A IgG, and

by —●— the values are shown of anti-*N. meningitidis* group C IgG.

Figure 3: In this figure, the chart is shown which is obtained for the determination of the bactericidal activity of the rabbit antisera on living strains of *N. meningitidis*.

On the abscissae the time (days) and on the ordinates the reciprocal of the logarithm of the dilu-

tion are reported,

and by $\square \text{---} \square$ the anti-group C bactericidal activity is shown.

Figure 4: the chart is shown which is obtained for the determination of the anti-diphtherin in antisera of Guinea-pigs immunized by the glycoprotein conjugate having trivalent immunogenic activity.

On the abscissae the time (days) and on the ordinates the concentrations, expressed as S.I. units/ml are reported.

By $\Delta \text{---} \Delta$ the anti-diphtherin is indicated.

Each point represents the geometrical average of the titrations obtained for each group of 10 animals.

EXAMPLE 1

Preparation of the Oligosaccharidic Haptens Derived from the Capsular Polysaccharide of *Streptococcus pneumoniae* Type 6A

Two mg of capsular polysaccharide of *S. pneumoniae* Type 6A is dissolved in 1 ml of aqueous solution containing 10 mM of acetic acid at pH 3.4.

The reaction of hydrolysis is carried out by maintaining the solution inside sealed ampuls dipped in oil bath, at a temperature of 100°C, for a time of 39 hours.

At the end of said time, the so-obtained oligosaccharides are separated from the reaction mixture, by chromatography over Sephadex G 15 - (Pharmacia, Uppsala), conditioned with a 200 mM solution of NaCl at pH 7.0, by operating at 4°C.

The chromatographic effluents are then analyzed according to the procedures as reported by Kabat in Exp. Immunochimistry, Ed. E.A. Rabat & Mayer, pages 538-541, 1964), Chen et al. (Anal. Chem. **28**, 1756-1758, 1956) and by Porro et al. - (Anal. Biochem. **118**, 301-306, 1981) to the purpose of verifying the presence of methyl-pentoses, phosphorus and reducing groups.

The results obtained show a phosphorus/rhamnose molecular ratio of 1 rhamnose/reducing groups of 3.

Upon analysis over Sephadex G-50, said oligosaccharides show a distribution constant (Kd) of 0.48, corresponding to a molecular weight (MW) of about 1500.

Upon N.M.R. analysis, the oligosaccharides result formed by about 8 monosaccharidic residues, among which the galactose residue, which is the immunodominant sugar, is found.

The N.M.R. analysis is typically carried out by dissolving 50 mg of the oligosaccharide in 1 ml of deuterium oxide (D_2O), using 5-mm probes at 35°C, a Varian Spectrometer model CFT 20, and operating at 20 Megahertz (20 MHz), using, as the inner standard, methanol, which has a $\delta = 39.5$ parts per million (ppm).

EXAMPLE 2

Preparation of the Oligosaccharidic Haptens Derived from the Capsular Polysaccharide of *Neisseria meningitidis* Group C

The test is carried out as in preceding Example 1, but using, for the hydrolysis of the capsular polysaccharide of *N. meningitidis* group C, a 0.5 mM solution of acetic acid, at pH = 5, for a time of 8 hours.

The oligosaccharidic haptens so obtained show, over Sephadex G-50, a Kd of 0.46, corresponding to a molecular weight of about 1500.

The chemical analysis of the chromatographic effluents shows the presence of terminal reducing groups, and of a residue of N-acetyl-neuraminic acid, also known as sialic acid.

Upon N.M.R. analysis, said haptens result constituted by 10-12 monosaccharidic residues, among which the immunodominant residue N-acetyl-neuraminic acid.

EXAMPLE 3

Determination of the Immunochemical Properties of the Oligosaccharidic Haptens Derived from the Capsular Polysaccharides of *S. pneumoniae* 8A and *N. meningitidis* group C.

The test is carried out as described by Porro et al., in Molec. Immunol., in press 1985.

Used are plastic plates for immunoelectrophoresis, subdivided into three compartments, A, B and C compartments, each containing 2 ml of 0.1% (w/v) agarose (Agarose M-LKB, Bromma).

Into compartment A, 0.05% (v/v) is introduced of rabbit antiserum, detector for the capsular polysaccharide of *S. pneumoniae* Type 6A, or for the capsular polysaccharide of *N. meningitidis* group C (Staten Seruminstitut, Copenhagen).

Into compartment B, 0.05% (v/v) is introduced of rabbit antiserum, detector for the capsular polysaccharidic antigen of *S. pneumoniae* Type 6A or of the capsular polysaccharidic antigen of *N. meningitidis* group C, previously incubated at 37°C

for 15 minutes, with respectively 5 nanograms of capsular polysaccharide of *S. pneumoniae* Type 6A and 5 nanograms of capsular polysaccharide *N. meningitidis* group C.

Into compartment C, 0.05% (v/v) is introduced of rabbit antiserum, detector for the capsular polysaccharide of *S. pneumoniae* Type 6A, or of rabbit antiserum, detector for the capsular polysaccharide *N. meningitidis* group C., previously incubated at 37°C for 15 minutes with 5, 50, 500 and 5000 nanograms of the corresponding oligosaccharidic haptens.

The capsular polysaccharides Type 6A and group C used as reference are diluted four times, each time by a 1:2 ratio (2, 4, 8, 16 total µg), and 10 µl of each dilution are introduced into each one of the 4 wells provided in agarose of Compartments B and C.

The plates so prepared are submitted to electrophoresis, operating at 70 V/cm, for a time period of from 1 to 2 hours. At the end, the plates are dried at 37°C, and the immunoprecipitated rockets are determined and quantified.

The sensitivity of this method is of about 20 ng/ml in the determination of the capsular polysaccharide of *S. pneumoniae* Type 6A, and of about 50 ng/ml in the determination of polysaccharide of *N. meningitidis* group C.

The inhibition of the oligosaccharidic haptens was evidenced when the immunoprecipitated rockets were higher in agarose compartment C, which contains the reference antiserum preincubated with the oligosaccharidic haptens, than in compartment B, containing the reference antiserum preincubated with the polysaccharidic antigens.

The minimum inhibiting concentration of the oligosaccharidic haptens obtained as reported in Examples 1 and 2, and their immunochemical specificity, expressed as the amount by weight of the haptens, to the antigens recognized by the reference rabbit antibodies is then determined according to as described by Porro et al. in *Molec. Immunol.*, in press, 1985.

In particular, it has been found that, for the oligosaccharidic haptens derived from the capsular polysaccharide of *S. pneumoniae* type 6A, the lower specificity value indicated the presence, in the structure, of the immunodominant sugar galactose, and for the oligosaccharidic haptens derived from the capsular polysaccharide *N. meningitidis*, the presence of the immunodominant acid: N-acetylneuraminic acid.

The specificity of the oligosaccharidic haptens, when compared to that of the respective polysaccharidic antigens was of respectively 1.08×10^{-3} for the oligosaccharidic hapten derived from the cap-

sular polysaccharide of *S. pneumoniae* Type 6A and 1.06×10^{-3} for the oligosaccharidic hapten derived from the polysaccharide of *N. meningitidis* group C.

EXAMPLE 4

Preparation of Protein CRM 197

CRM 197, produced by the strain *Corynebacterium diphtheriae* C7 ($\beta^{tox-197}$), is separated from the culture medium by molecular filtration, by using a Millipore XM-50 (NMWL 5×10^{-4}) membrane.

The protein is then precipitated, by adding to the filtrate a solution of ammonium sulphate up to 65% of saturation.

The precipitated protein is then separated by centrifugation, and dissolved in 0.01 M phosphate buffer (pH 7.2).

Protein CRM 197 is subsequently purified by ion-exchange chromatography.

Used is a 2.5×100 cm DEAE Sepharose 6B/CL column (Pharmacia, Uppsala), conditioned in 0.01 M phosphate buffer at pH 7.2, and as the eluent a buffer solution with a 0.09 M concentration of NaCl.

80% of so-obtained CRM 197 is in its native molecular form, as determined by sodium dodecyl sulphate-Electrophoresis over polyacrylamide gel - (SDS -PAGE) under reducing conditions, according to as reported by Pappenheimer et al. (*Immunochemistry* 9, 891-906, 1972).

The purity of the protein is of about 400 flocculation limit (Lf)/mg.

EXAMPLE 5

Activation of the Oligosaccharidic Haptens Derived from the Capsular Polysaccharide of *S. Pneumoniae* Type 6A and from the Capsular Polysaccharide *N. meningitidis* group 6C and Their Conjugation with CRM 197

a) Introduction of the primary amino groups into the terminal reducing groups of the oligosaccharidic haptens.

The oligosaccharidic haptens, obtained as reported in Examples 1 and 2, are separately dissolved in distilled water, to an end concentration of 10 mg/ml. To each solution, 50 mg/ml of ammonium chloride, and 20 mg/ml of NaBH_4CN are then added.

The solution containing the oligosaccharidic haptens derived from capsular polysaccharide *S. pneumoniae* Type 6A is adjusted to pH 9 by 1 N NaOH and is kept, inside sealed ampuls, under

mild stirring, at a temperature of 50°C over two weeks. The solution containing the oligosaccharidic haptens derived from the capsular polysaccharide of *N. meningitidis* group C is adjusted to pH 8 and is kept, inside sealed ampuls, under mild stirring, at a temperature of 37°C for one week.

At the end of the reaction, the oligosaccharidic haptens are purified by gel-chromatography, using a Sephadex G15 column conditioned in buffer solution with mM NaCl at pH 7.0.

The chromatographic effluents having $K_d \leq 0.1$ and a chemical activity for respectively methyl pentose groups, phosphorus and amino groups, or N-acetyl-neuraminic acid and amino groups are collected and combined.

b) Conversion of amino-group-hapten derivatives into the corresponding monofunctional esters

The oligosaccharidic haptens into which the amino groups have been introduced are diluted to a concentration of 10 mg/ml in an aqueous solution of dimethylsulphoxide (at 80-95% v/v).

Said solutions are added dropwise, over a time ranging from 1 to 2 hours, to dimethylsulphoxide containing di-succinimidyl ester of adipic acid - (DEA) in molar excess relatively to the amino groups introduced into the oligosaccharidic haptens (10:1).

The esterification reaction is carried out by keeping the solutions at 20-25°C for a time period of 4 hours. At the end of said time period, the esterified oligosaccharidic haptens are precipitated by the addition of 1,4-dioxane to an end concentration of 90% (v/v), and are subsequently separated from the reaction mixture by centrifugation.

The conversion of the amino groups into the corresponding active esters is then verified by the method as reported by Miron-Wilchek (Anal. Biochem. 126, 433-435, 1982).

c) Conjugation of the esterified oligosaccharidic haptens to CRM 197

To an aqueous solution containing 2 mg/ml of CRM197, a solution is added of dimethylsulphoxide containing the activated oligosaccharidic hapten derived from the capsular polysaccharide of *S. pneumoniae* Type 6A in 10:1 molar ratio to the protein, until an end concentration of dimethylsulphoxide in water of 20% (v/v) is obtained.

The mixture so obtained is kept, under mild stirring, at room temperature (20-25°C) for 15 hours. At the end of said time period, the organic solution containing the activated oligosaccharidic hapten derived from the capsular polysaccharide of *N. meningitidis* group C is added in a 10:1 molar ratio to the protein, and the resulting mixture is kept 15 hours at room temperature.

According to an alternative route, the conjugation reaction can be carried out by adding, at the same time, to the CRM-197-containing solution, the two activated oligosaccharidic haptens respectively derived from the capsular polysaccharide of *S. pneumoniae* Type 6A and from the capsular polysaccharide of *N. meningitidis* group C.

The glycoprotein conjugates so obtained are separated by gel-chromatography, using a Sephadex G-100 column (Pharmacia, Uppsala), conditioned in 0.01 M phosphate buffer containing 0.2 M NaCl (pH = 7.0).

The eluted fractions showing a chemical activity for the proteins, phosphorus and sialic acid are combined and filtered through a millipore membrane having a porosity of 0.22 μm .

The chemical analysis of the glycoprotein conjugate so obtained shows a content of protein of 87% (w/w), of Type-6A oligosaccharidic hapten of 8% (w/w), and of group-C oligosaccharidic hapten of 5% (w/w).

These values, converted into mol, show a degree of substitution, per each mol of protein CRM 197, of 4 mol of oligosaccharidic haptens of Type 6A, and of 2 mol of oligosaccharidic haptens of group C.

The physical-chemical characterization of the glycoprotein conjugate has been determined by SDS-PAGE and a gradient of acrylamide of 3-9%, according to Laemmli's procedure (Nature 227, 680-685, 1970) (Fig. 1).

The immunochemical specificity of monoclonal antibodies for different epitopes of protein CRM 197, as compared to that observed for the glycoprotein conjugate wherein the protein is CRM 197, is reported in Table 1.

The values, obtained from three experimental tests, are determined by using the ELISA methodology.

The specificity percentage of the monoclonal antibodies for the protein conjugate was computed assuming a value of 100 for the specificity observed by measuring the absorbance values, in the ELISA procedure, for CRM 197.

I a b l e 1

Monoclonal Antibodies	Recognized Epitopes in CRM-127	Number of Residual Aminoacids in the Epitopes		% Specificity for	
		Lys	Arg	CRM-127	Glycoproteinic conjugate
Group I					
(Clone 6 B 12)	1-156	14	2	100	31
Group II b					
(Clone 2 A 5)	157-193	2	5	100	<10
Group III a					
(Clone 2 A 7)	293-345	1	-	100	21
Group IV C I)					
(Clone 2 E 10)	465-535	6	2	100	95

As it can be seen from the preceding Table, the most important epitope of CRM 197, localized in the 465-535 sequence of the protein, is present in the protein CRM 197 of the glycoprotein conjugate.

This epitope is the most important immunodeterminant of the diphtherin, it being able to induce neutralizing antibodies by the inhibition of the cellular bond of the same toxin on sensible cells.

EXAMPLE 5

Immunological Characteristics of Animal Sera Obtained After Immunization by the Glycoprotein Conjugate Having Trivalent Immunogenic Activity Obtained in Example 5

Two groups of 10 rabbits of weight of about 2 kg, and one group of 10 Guinea-pigs, of weight of 350 g, are inoculated by subcutaneous way with two doses of the glycoprotein conjugate absorbed on a support of AIPO₄ (37.5 mg of glycoprotein conjugate/1.5 mg of AIPO₄).

Each dosis contains 12.5 Lf of CRM 197, 4.8 ng of oligosaccharidic hapten derived from the capsular polysaccharide of *S. pneumoniae* Type 6a, and 3 ng of oligosaccharidic hapten derived from the capsular polysaccharide of *N. meningitidis* group C.

The second dosis is inoculated 28 days after the 1st dosis. Serum collections are carried out on the rabbits soon after the administration of the two doses, and 11 days after the administration of the second dosis.

Rabbit antisera are analyzed by an immunoenzymatic method (ELISA) according to A. Voller et al., "Microplate Enzyme-Linked Immunosorbent Essay for Chagas Disease", *Lancet* **1**, 426 (1975), to the purpose of determining the presence of IgG immunoglobulins specific to the oligosaccharidic hapten derived from the capsular polysaccharide of *S. pneumoniae* Type 6A and to the oligosaccharidic hapten derived from the capsular polysaccharide of *N. meningitidis* group C (Fig. 2).

The bactericidal activity of rabbits antisera has been verified on living strains of *N. meningitidis* according to the method as described by Wong et al. (*J. Biol. Stand.* **5**, 197-215, 1977) and the results have confirmed the development of a meaningful bactericidal activity soon after the administration of each of the doses of glycoprotein conjugate (Fig. 3).

The antisera from Guinea-pigs are analyzed for determining the content of anti-diphtherin, according to the U.S. Pharmacopoeia method.

A horse antiserum for diphtherin is used as the standard in the titration of the diphtherin used in the biological analysis, and in Fig. 4, the chart obtained is illustrated.

In all the animals, before the immunization, the absence of specific antibodies has been determined.

The immunization of control animals with only oligosaccharidic haptens or capsular polysaccharides had shown the absence of immunogenicity of said purified compounds.

Claims

1. Glycoprotein conjugates having trivalent immunogenic activity, obtained by binding, by a covalent bond, to a proteinic antigen selected among CRM 197, tetanus toxoid, or pertussis toxin, at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, after the preliminary activation of the said oligosaccharidic haptens by the introduction of terminal ester groups.

2. Glycoprotein conjugates having trivalent immunogenic activity according to claim 1, wherein the oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium is selected among those deriving from bacteria belonging to the group of *Streptococcus pneumoniae* and of *Streptococcus β-hemolyticus* and the oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium is selected among those deriving from bacteria belonging to the group of *Neisseria meningitidis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa* and *Escherichia coli*.

3. Glycoprotein conjugate having trivalent immunogenic activity according to claim 1, wherein the protein is CRM 197, the oligosaccharidic hapten derived from the capsular polysaccharide of the gram-positive bacterium is obtained from *Streptococcus pneumoniae* and the oligosaccharidic hapten derived from the capsular polysaccharide of the gram-negative bacterium is obtained from *Neisseria meningitidis*, with a molecular ratio between the three components of 1/4/2.

4. Glycoprotein conjugates having trivalent immunogenic activity according to claims from 1 to 4, wherein the activated oligosaccharidic haptens have a molecular weight of about 1500.

5. Process for the preparation of glycoprotein conjugates having trivalent immunogenic activity according to claims from 1 to 4, characterized in that to a proteinic antigen selected among CRM 197, tetanus toxoid, or pertussis toxin, at least one

oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium, and at least one oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium, are either contemporaneously or sequentially bound, by a covalent bond, after the preliminary activation of the said oligosaccharidic haptens by the introduction of terminal ester groups.

6. Process according to claim 5, characterized in that the oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium is obtained from bacteria belonging to the group of *Streptococcus pneumoniae* or of *Streptococcus β -hemolyticus* and the oligosaccharidic hapten derived from the capsular polysaccharide of a gram-negative bacterium is obtained from bacteria belonging to the group of *Neisseria meningitidis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa* and *Escherichia coli*.

7. Process according to claim 6, characterized in that the protein is CRM 197, the oligosaccharidic hapten derived from the capsular polysaccharide of a gram-positive bacterium is obtained from *Streptococcus pneumoniae*, and the oligosaccharidic

hapten derived from the capsular polysaccharide of the gram-negative bacterium is obtained from *Neisseria meningitidis*.

8. Process according to claims from 5 to 7, characterized in that the conjugation reaction between the proteinic antigen and the activated oligosaccharidic haptens is carried out, either contemporaneously or sequentially, in a liquid medium, in the presence of dimethylsulphoxide, at a temperature comprised within the range of from 20°C to 25°C and for a corresponding time period of from 4 to 24 hours.

9. Vaccine against capsulate gram-negative and gram-positive bacteria, which comprises a therapeutically effective amount of a glycoprotein conjugate having trivalent immunogenic activity, as defined in claims from 1 to 4, and a pharmacologically acceptable amount of a carrier selected among those known from the art.

10. Method for inducing an active immunization in mammals, against infections caused by capsulate gram-positive and gram-negative bacteria, which comprises the administration of a vaccine containing an immunogenic amount of the glycoprotein conjugate having trivalent immunogenic activity as defined in claims from 1 to 4.

30

35


40




45

50

55

10

Fig.1

Fig. 2Fig. 3Fig. 4